

6.1 VOLUMES USING CROSS-SECTIONS

The volume of any solid with cross-sectional area $A(x)$ from $x = a$ to $x = b$ is

$$V = \int_a^b A(x)dx \quad (1)$$

Disks.

Specifically, for rotations using disks (i.e. regions without gaps) about the x and y axes are respectively,

$$V = \int_a^b \pi R(x)^2 dx \quad (2)$$

$$V = \int_a^b \pi R(y)^2 dy \quad (3)$$

- (1) Find the volume of a sphere centered at the origin with radius r .
Solution: Consider circular cross-sections, i.e. circles cutting the sphere perpendicular to the x-axis of radius y . Then, by the distance formula (or Pythagorean theorem), $R(x)^2 = y^2 = r^2 - x^2$. So,

$$V = \int_a^b \pi R(x)^2 dx = \int_{-r}^r \pi(r^2 - x^2)dx = \frac{4}{3}\pi r^3.$$

- (2) Find the volume of a region bounded by $y = \sqrt{x}$, $x = 0$, and $x = 1$ about the x-axis.

$$\text{Solution : } R(x) = \sqrt{x} \Rightarrow V = \int_0^1 \pi x dx = \frac{\pi}{2}$$

- (3) Find the volume of a region bounded by $y = x^3$, $x = 0$, and $y = 8$ about the y-axis.

Solution: Since we are revolving around the y-axis we need to solve for x as a function of y , i.e. $x = y^{1/3}$, then

$$R(y) = y^{1/3} \Rightarrow V = \int_0^8 \pi y^{2/3} dy = \frac{96}{5}\pi$$

Washers.

For rotations using washers (i.e regions with gaps) about the x and y axes are respectively,

$$V = \int_a^b \pi[R(x)^2 - r(x)^2]dx \quad (4)$$

$$V = \int_a^b \pi[R(y)^2 - r(y)^2]dy \quad (5)$$

Where R is the radius of the larger region, and r is the radius of the smaller region (i.e. gap).

- (1) Find the volume of the region between $y = x$ and $y = x^2$ revolved about the x-axis.

Solution: Since $y = x$ is on top, it sweeps out a larger region than $y = x^2$, so $R(x) = x$ and $r(x) = x^2$. Then,

$$V = \int_a^b \pi[R(x)^2 - r(x)^2]dx = \int_0^1 \pi[x^2 - x^4]dx = \frac{2\pi}{15}$$

- (2) What if we revolve it around $y = 2$?

Solution: Since we are revolving about an axis above our region, the function on the bottom will now sweep out a larger region. This can be seen by sketching the curves. So, $R(x) = 2 - x^2$ and $r(x) = 2 - x$. Then,

$$V = \int_a^b \pi[R(x)^2 - r(x)^2]dx = \int_0^1 \pi[(2 - x^2)^2 - (2 - x)^2]dx = \frac{8\pi}{15}$$

- (3) Now lets revolve it around $x = -1$.

Solution: First we must solve for x as a function of y now. We get, $x = y$ and $x = \sqrt{y}$, respectively. Since we are revolving about an axis to the left of our region, the function on the right will sweep out a larger region. So, $R(y) = \sqrt{y} - (-1)$ and $r(y) = y - (-1)$. Then,

$$V = \int_a^b \pi[R(y)^2 - r(y)^2]dy = \int_0^1 \pi[(1 + \sqrt{y})^2 - (1 + y)^2]dy = \frac{\pi}{2}$$

Tougher examples.

- (1) Find the volume of a region such that the base is a circle centered at the origin of radius $r = 1$ and cross-sections perpendicular to the x-axis that are equilateral triangles.

Solution: Notice the base of the triangle will be $2y$, and since they are equilateral triangles we can split the triangle in half to get a “30, 60, 90” triangle with base y , so the height of the triangles will be $\sqrt{3}y$. This gives us a cross-sectional area of $A(y) = \sqrt{3}y^2$, but we have cross-sections perpendicular to the x-axis, so we need $A(x)$. Notice a circle is given by the equation $y^2 + x^2 = r^2$, but $r = 1$, so $y^2 = 1 - x^2$. Hence, $A(x) = \sqrt{3}(1 - x^2)$. Then,

$$V = \int_{-1}^1 \sqrt{3}(1 - x^2)dx = \frac{4\sqrt{3}}{3}.$$

- (2) Find the volume of a square pyramid such that the sides of the square are length L and height h .

Solution: There are many ways to do this, but perhaps the easiest is to consider a triangle with the head at the origin and base at $x = L$ with square cross-sections perpendicular to the x-axis. So, at an arbitrary x , the cross-section is a square of length, say ℓ . Notice, the entire triangle and a triangle at any arbitrary x will be similar triangles, and hence have the same ratios. Therefore, $\frac{x}{h} = \frac{\ell}{L}$, i.e. the ratio of the heights equal the ratio of the lengths. This gives $\ell = \frac{L}{h}x$. Then, $A(x) = (Lx/h)^2$, and

$$V = \int_0^h \frac{L^2}{h^2}x^2dx = \frac{1}{3}L^2h$$

- (3) Find the volume of a region such that the base is a semicircle centered at the origin with radius $r = 4$ and cross-sections perpendicular to the x-axis that are “30, 60, 90” triangles - 30° with respect to the x-axis.

Solution: Notice the base at any arbitrary x will be of length y . The height of the triangle will be $y/\sqrt{3}$, then $A(y) = \frac{1}{2\sqrt{3}}y^2$. Since $r = 4$, $y^2 = 16 - x^2$. Then, $A(x) = \frac{1}{2\sqrt{3}}(16 - x^2)$. Then,

$$V = \int_{-4}^4 \frac{1}{2\sqrt{3}}(16 - x^2)dx = \frac{128}{3\sqrt{3}}.$$

6.2 CYLINDRICAL SHELLS

Another method to do volumes of revolutions is through cylindrical shells. This method is a lot less intuitive, and hence requires more practice. I wonder if any of you guys are actually reading this... Basically think of infinitesimal cylinders filling up a region. We know the area of the side of the cylinder is $A = 2\pi rh$. So, by summing up these infinitesimal cylinders we get the following formulas for rotation about the y-axis and x-axis respectively,

$$V = \int_a^b 2\pi x f(x) dx \quad (6)$$

$$V = \int_a^b 2\pi y f(y) dy \quad (7)$$

- (1) Find the volume of the region bounded by $y = 2x^2 - x^3$ and $y = 0$ revolved about the y-axis.

Solution: Here the radius of each cylinder will be $r = x$ and the height will be $h = y = 2x^2 - x^3$. Then,

$$V = \int_a^b 2\pi x f(x) dx = \int_0^2 2\pi x(2x^2 - x^3) dx = \frac{16}{5}\pi$$

- (2) Find the volume of the region bounded by $y = x$ and $y = x^2$ revolved about the y-axis.

Solution: The radius is $r = x$ and the “height” is $h = x - x^2$, then

$$V = \int_a^b 2\pi x f(x) dx = \int_0^1 2\pi x(x - x^2) dx = \frac{\pi}{6}$$

- (3) Find the volume of the region bounded by $y = \sqrt{x}$, $x = 0$, and $x = 1$ revolved about the x-axis.

Solution: The radius is $r = y$ and the “height” is $h = 1 - y^2$, then

$$V = \int_a^b 2\pi y f(y) dy = \int_0^1 2\pi y(1 - y^2) dy = \frac{\pi}{2}$$

- (4) Find the volume of a region bounded by $y = x - x^2$ and $y = 0$ revolved about the line $x = 2$.

Solution: Since our axis of revolution is towards the right, the radius of our cylinders will be $r = 2 - x$ and the height is $h = y = x - x^2$. Hence, our area is $A(x) = 2\pi(2 - x)(x - x^2)$. Then,

$$V = \int_0^1 2\pi(2 - x)(x - x^2) dx = \frac{\pi}{2}$$